



The Meeting of the Americas

You may print by clicking on this [Print](#) button. To return to the previous page, close this browser window or click the 'X' button in the top right corner of the page.

ID# P31B-04

Location: Rafain Expocenter II-III Exhibit Hall

Time of Presentation: Aug 11 8:00 AM - 12:30 PM

Present-day and early heat flows on Mars: implications for thermal evolution

*J. Ruiz*¹; *V. Lopez*²

1. Departamento de Geodinamica, Universidad Complutense de Madrid, Madrid, Spain.

2. Instituto de Geologia Economica, CSIC-UCM, Madrid, Spain.

There is currently a very interesting and productive debate on the present-day heat flow and thermal state of Mars. The very high (>300 km) effective elastic thickness deduced for the north polar region from the non significant flexure caused by loading due to the north pole cap, is related to a low surface heat flow at the present-time, hardly reconcilable with most of thermal history models. This low heat flow could be indicative of sub-chondritic heat-producing elements (HPE) abundances, a limited influence of secular cooling, or simply a regional variability of surface heat flow. On the other hand, effective elastic thicknesses of Noachian-loaded regions are very low (typically ~15 km or lower), indicating high heat flows, which is espectral for a young planet. Otherwise, the depth to the brittle-ductile transition beneath Late Noachian-Early Hesperian lobate scarps (considered to be related with large thrust faults) is ~20-35 km, which can in turn be converted to heat flows of ~25-45 mW m⁻². These values are lower than those predicted by thermal history models of Mars, and than the heat flow equivalent to the total radioactive heat generation. The discrepancy between deduced and predicted Late Noachian-Early Hesperian lobate heat flows could be explained by a limited contribution from secular cooling, as already proposed for explain the present-day heat flow in the north polar region. However, it could also be explained an efficient hydrothermal cooling of the upper crust, which was previously proposed to explain the preservation of large-scale crustal thickness variations observed in the southern highlands. Thus, the comparison between heat flows estimated from both the effective elastic thicknesses of the lithosphere and the depth to the brittle-ductile transition, and those predicted by thermal history models, may serve to put constraints on the evolution of the efficiency of several ways of heat loss (including vigorous hydrothermal coolings) on Mars.

Contact Information

Javier Ruiz, Madrid, Spain, 28040, [click here](#) to send an email

ScholarOne Abstracts® (patent #7,257,767 and #7,263,655). © [ScholarOne](#), Inc., 2010. All Rights Reserved.

ScholarOne Abstracts and ScholarOne are registered trademarks of ScholarOne, Inc.

[Terms and Conditions of Use](#)